

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

DIRECTIONS FOR USE

Do not apply to food or feed crops. Do not use where stored food or feed (such as grain or meat) might become contaminated. Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

Applicators shall wear gloves and use a scoop when mixing, applying and broadcasting this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Storage:** Store only in original container in a cool, dry location inaccessible to children and pets. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or insect infestation.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal:** Fiber Drums with Liners: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in sanitary landfill or by incineration if allowed by State and Local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

RESTRICTED USE PESTICIDE DUE TO ACUTE AVIAN TOXICITY

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the certified Applicator's Certification.

24(c) Supplemental Label

For use only in the State of Nevada. When controlling Common Ravens a USFWS depredation permit must be obtained.

AVITROL

DOUBLE STRENGTH WHOLE CORN

Active Ingredient: 4-aminopyridine	1.0%
Inert Ingredients:	99.0%
Total:	100.0%

Avitrol is a poison with flock alarming properties used for the control of Black-billed Magpies (*Pica hudsonia*) and Common Ravens (*Corvus corax*) in, on or in the area of structures, feeding, nesting and roosting sites, in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid

Have label with you when obtaining treatment advice.

IF SWALLOWED: -Call a poison control center, doctor, or 800-424-9300 immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to do so by the poison control center or doctor.

IN EYES: -Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center, doctor, or 800-424-9300 immediately for treatment advice.

Manufactured By
AVITROL CORPORATION
7644 East 46th Street
Tulsa, OK 74145-6370
(918) 622-7763 * (800) 633-5069
EPA Reg. No. 11649-8
SLN NV-130002

SPECIMEN LABEL

PREBAITING DIRECTIONS

This product must not be applied where non-target birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no nontarget birds are feeding on the prebait. Prebaiting with untreated grains of the same composition as the Avitrol carrier is usually essential to the effective deterrence of birds with Avitrol.

DILUTION DIRECTIONS

Avitrol treated grain should be thoroughly mixed with untreated grain of the same composition as the Avitrol carrier at the dilution ratios indicated below. For the control of Black-billed Magpies and Common Ravens with Avitrol, no dilution ratio less than one part treated to nine parts untreated bait is recommended.

BAITING DIRECTIONS

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necessary number of target birds. After the birds' feeding pattern has been established through prebaiting, replace untreated bait with diluted treated bait only at sites where the target birds are actively feeding. Do not apply treated bait to inactive feeding sites. Where uneaten bait may be a hazard to other birds or animals, it must be picked up at the end of each day. When this product is used in or around occupied buildings, pick up and dispose of dead birds by burial. This product must not be applied where nontarget bird species are feeding. During the first few days, baiting with Avitrol may require successive applications until control is obtained. Note that retreatment may also be necessary after rainfall.

Notice: Buyer and/or user assumes all risks of use and or handling of these materials contrary to label instructions.